

SECOND SEMESTER DIPLOMA EXAMINATION IN ENGINEERING/  
TECHNOLIGY- OCTOBER, 2011

TECHNICAL MATHEMATICS- II  
(Common – Except DCP and CABM)

[Time: 3 hours

(Maximum marks: 100)

Marks

PART –A  
(Maximum marks: 10)

(Answer all questions. Each question carries 2 marks)

I.

(a) Evaluate  $\lim_{\theta \rightarrow 0} \frac{\sin \theta}{6\theta}$

We have  $\lim_{\theta \rightarrow 0} \frac{\sin m\theta}{n\theta} = \frac{m}{n}$

$\therefore \lim_{\theta \rightarrow 0} \frac{\sin \theta}{6\theta} = \frac{5}{6}$

(b) If  $y = \sin^2 x$ , find  $\frac{d}{dx}$

$y = \sin^2 x$

$\therefore \frac{d}{dx} = 2\sin x \cos x$

(c) For what value of 'x' will be the tangent to the curve  $y = \frac{x}{x^2+1}$

$y = \frac{x}{x^2+1}$

$\frac{dy}{dx} = \frac{(x^2+1)x - x \times 2x}{(x^2+1)^2}$

(Apply Quotient Rule)

$$= \frac{(x^2+1) - 2x^2}{(x^2+1)^2}$$

$$\frac{dy}{dx} = \frac{1-x^2}{(1+x^2)^2}$$

Since the tangent is parallel to x – axis

$$\therefore \frac{d}{dx} = 0$$

$$\text{ie } \frac{1-x^2}{(1+x^2)^2} = 0$$

$$\text{ie } 1 - x^2 = 0$$

$$x^2 = 1$$

$$\text{ie } x = \pm 1$$

(d) Find  $\int \cot x \, dx$

$$\int \cot x \, dx = \int \frac{\cos x}{\sin x} \, dx$$

Put  $\sin x = u$

$$\frac{du}{dx} = \cos x$$

$$du = \cos x \, dx$$

$$\therefore \cos x \, dx = du$$

$$= \int \frac{du}{u}$$

$$= \int \frac{1}{u} \, du$$

$$= \log u$$

$$= \log \sin x + c$$

(e) Solve  $\int \frac{d\theta}{\theta} = \frac{5}{2} \theta$

Apply variable – separable form

$$\int \frac{d\theta}{\theta} = \frac{5}{2} dt$$

Integrating on both sides

$$\int \frac{d\theta}{\theta} = \int \frac{5}{2} dt$$

$$\log \theta = \frac{5}{2}t + c$$

PART -B

Answer any five questions. Each question carries 6 marks

II.

(a) Evaluate

i.  $\lim_{x \rightarrow -2} \frac{x^3 + 8}{x + 2}$

$$\lim_{x \rightarrow -2} \frac{x^3 + 8}{x + 2}$$

Here n = 3, a = -2

$$\lim_{x \rightarrow a} \frac{x^n - a^n}{x - a} = n \times a^{n-1}$$

$$\begin{aligned} \therefore \lim_{x \rightarrow -2} \frac{x^3 + 8}{x + 2} &= 3 \times (-2)^{3-1} \\ &= 3 \times (-2)^2 \\ &= 3 \times 4 \\ &= 12 \end{aligned}$$

ii. Examine whether the function given by  $f(x) = \begin{cases} x^2 + 3, & \text{if } x \neq 0 \\ 1, & \text{if } x = 0 \end{cases}$  is continuous at  $x=0$

$$f(x) = \begin{cases} x^2 + 3, & \text{if } x \neq 0 \\ 1, & \text{if } x = 0 \end{cases}$$

$$f(0) = 1$$

$$\lim_{x \rightarrow 0} f(x) = \lim_{x \rightarrow 0} x^2 + 3$$

$$= 0^2 + 3$$

$$= 3$$

$$\therefore \lim_{x \rightarrow 0} f(x) \neq f(0)$$

$\therefore$  Function is not continuous at  $x = 0$

(b) Find  $\frac{dy}{dx}$  if

i.  $y = (1 - 2x + 7x^2)^{10}$

$$y = (1 - 2x + 7x^2)^{10}$$

$$\frac{dy}{dx} = 10(1 - 2x + 7x^2)^9 \cdot (-2 + 14x)$$

ii.  $y = \log(\operatorname{cosec}x - \cot x)$

$$y = \log(\operatorname{cosec}x - \cot x)$$

$$\begin{aligned} \frac{dy}{dx} &= \frac{1}{(\operatorname{cosec}x - \cot x)} \cdot (-\operatorname{cosec}x \cdot \cot x + \operatorname{cosec}^2 x) \\ &= \operatorname{cosec}x \frac{\operatorname{cosec}x - \cot x}{\operatorname{cosec}x - \cot x} \\ &= \operatorname{cosec}x \end{aligned}$$

(c) The distance 's' meters travelled by a particle is given by,  $s = ae^{2t} + be^{-2t}$ , where, 't' represents the time. Prove that the acceleration varies as the distance.

$$s = ae^{2t} + be^{-2t}$$

$$\begin{aligned} \frac{ds}{dt} &= ae^{2t} \cdot 2 + be^{-2t} \cdot (-2) \\ &= 2ae^{2t} - 2be^{-2t} \end{aligned}$$

$$\begin{aligned} \frac{d^2s}{dt^2} &= 4ae^{2t} + 4be^{-2t} \\ &= 4(ae^{2t} + be^{-2t}) \\ &= 4 \times s \end{aligned}$$

$$\frac{d^2s}{dt^2} = k \times s, \text{ where } k = 4$$

Acceleration varies as the distance

(d) The bending moment of a rod 10m long and weighing 40kg and resting on support supports at its ends at a distance of 'x' meters from one end is given by  $M = 2(10x - x^2)$ . Find the maximum bending moment?

$$M = 2(10x - x^2) = 20x - 2x^2$$

$$\frac{dM}{dx} = 20 - 4x$$

$$\frac{d^2m}{dx^2} = -4 < 0$$

$$\therefore \frac{dm}{dx} = 0 \implies 20 - 4x$$

$$20 = 4x$$

$$x = \frac{20}{4} = 5$$

$\therefore$  at  $x = 5$ ,  $m$  is maximum

$$\text{Maximum value, } m = 20x - 2x^2$$

$$= 20 \times 5 - 2 \times 5^2$$

$$= 100 - 50$$

$$= 50$$

(e) Integrate with respect to  $x$

i.  $\frac{1 + \sin x}{\cos^2 x}$

$$\frac{1 + \sin x}{\cos^2 x}$$

$$\therefore \int \frac{1 + \sin x}{\cos^2 x} dx$$

$$= \int \frac{1}{\cos^2 x} dx + \int \frac{\sin x}{\cos^2 x} dx$$

$$= \int \sec^2 x dx + \int \frac{\sin x}{\cos^2 x} \times \frac{1}{\cos x} dx$$

$$= \int \sec^2 x dx + \int \sec x \tan x dx$$

$$= \tan x + \sec x + c$$

ii.  $\cos^2 2x$

$$\text{We have } \cos^2 2x = \frac{1 + \cos 4x}{2}$$

$$\therefore \int \cos^2 2x dx = \int \frac{1 + \cos 4x}{2} dx$$

$$= \int \frac{1}{2} dx + \int \frac{\cos 4x}{2} dx$$

$$= \frac{1}{2} \int 1 dx + \int \frac{\cos 4x}{2} dx$$

$$= \frac{1}{2}x + \frac{1}{2} \frac{\sin 4x}{4} + c$$

(f) Find  $\int_0^2 x^2 \log x \, dx$

$$\int_0^2 x^2 \log x \, dx = \int_0^2 \log x \, x^2 \, dx$$

$$U = \log x$$

$$V = x^2$$

$$\int u v \, dx = u \int v \, dx - \int \left( \frac{du}{dx} \int v \, dx \right) dx$$

$$\int_0^2 \log x \, x^2 \, dx = \log x \frac{x^3}{3} - \int \left( \frac{x^3}{3} \cdot \frac{1}{x} \right) dx$$

$$= \log x \frac{x^3}{3} - \int \frac{x^2}{3} dx$$

$$= \frac{x^3}{3} \log x - \frac{x^3}{9} + c$$

$$\therefore \int_0^2 \log x \, x^2 \, dx = \left[ \frac{x^3}{3} \log x - \frac{x^3}{9} \right]$$

$$= \frac{8}{3} \log 2 - \frac{8}{3} - 0 - 0$$

$$= \frac{8}{3} \log 2 - \frac{8}{9}$$

(g) Solve  $(1 + x^2) \frac{dy}{dx} + y = e^{\tan^{-1} x}$

Dividing by  $(1 + x^2)$ , we get

$$\frac{dy}{dx} + \frac{y}{1+x^2} = \frac{e^{\tan^{-1} x}}{1+x^2}$$

This is of the form  $\frac{dy}{dx} + Py = Q$

$$\text{Here } p = \frac{1}{1+x^2} \quad Q = \frac{e^{\tan^{-1} x}}{1+x^2}$$

$$\therefore \text{I.F.} = e^{\int p \, dx} = e^{\int \frac{1}{1+x^2} \, dx} = e^{\tan^{-1} x}$$

$$\therefore \text{Solution is } Y \times \text{IF} = \int (Q \times \text{IF}) \, dx$$

$$Y \int x e^{\tan^{-1}x} = \int \left( \frac{e^{\tan^{-1}x}}{1+x^2} \times e^{\tan^{-1}x} \right) dx = \int u du \quad \text{where } u = e^{\tan^{-1}x}$$

$$= \frac{e^{2 \tan^{-1}x}}{2}$$

$$= \log u$$

$$Y \int x e^{\tan^{-1}x} = \log e^{\tan^{-1}x} + c$$

### PART -C

(Maximum mark: 60)

Answer four full questions. Each question carries 15 marks.

III.

(a) Evaluate  $\lim_{x \rightarrow 0} \frac{\sqrt{1+x}-1}{x}$

$$\lim_{x \rightarrow 0} \frac{\sqrt{1+x}-1}{x} = \lim_{x \rightarrow 0} \frac{(1+x)^{\frac{1}{2}} - 1^{\frac{1}{2}}}{(x+1)^{-1} - 1}$$

$$= n a^{n-1}$$

$$n = \frac{1}{2}$$

$$a = 1$$

$$\lim_{x \rightarrow 0} \frac{\sqrt{1+x}-1}{x} = \frac{1}{2} x^{-1} 1^{\frac{1}{2}-1}$$

$$= \frac{1}{2} x^{-1} 1$$

$$= \frac{1}{2}$$

(b)  $Y = \frac{x \tan^{-1}x}{1+x^2}$  find  $\frac{dy}{dx}$

$$Y = \frac{x \tan^{-1}x}{1+x^2}$$

$$\frac{dy}{dx} = \frac{(1+x^2) \left[ x \times \frac{1}{1+x^2} + \tan^{-1}x \times 1 \right] - x \tan^{-1}x \times 2x}{(1+x^2)^2}$$

$$= \frac{(1+x^2) \left[ \frac{x}{1+x^2} + \tan^{-1} x \right] - 2x^2 \tan^{-1} x}{(1+x^2)^2}$$

(c) Using 1<sup>st</sup> principle find the derivatives of cosx

$$Y = \cos x$$

$$f(x) = \cos x$$

$$f(x+\Delta x) = \cos(x+\Delta x)$$

$$\begin{aligned} \therefore \frac{dy}{dx} &= \lim_{x \rightarrow 0} \frac{f(x+\Delta x) - f(x)}{\Delta x} \\ &= \lim_{x \rightarrow 0} \frac{\cos(x+\Delta x) - \cos x}{\Delta x} \\ &= \lim_{x \rightarrow 0} \frac{-2 \sin\left(\frac{x+\Delta x+x}{2}\right) \sin\left(\frac{x+\Delta x-x}{2}\right)}{\Delta x} \\ &= \lim_{x \rightarrow 0} \frac{-2 \sin\left(\frac{2x+\Delta x}{2}\right) \sin\left(\frac{\Delta x}{2}\right)}{\Delta x} \\ &= \lim_{x \rightarrow 0} \frac{\sin\left(x+\frac{\Delta x}{2}\right) \sin\left(\frac{\Delta x}{2}\right)}{\frac{\Delta x}{2}} \\ &= \lim_{x \rightarrow 0} \sin\left(x+\frac{\Delta x}{2}\right) \times 1 \\ &= -\sin\left(x+\frac{0}{2}\right) \\ &= -\sin x \end{aligned}$$

(d) If  $y = \sin^{-1} x$ , prove that  $(1-x^2)y'' - xy' = 0$

$$Y = \frac{1}{\sqrt{1-x^2}}$$

$$Y' \frac{1}{\sqrt{1-x^2}} = 1$$

Differentiating on both sides

$$y' \frac{1}{2\sqrt{1-x^2}} x - 2x + \sqrt{1-x^2} \times y'' = 0$$



$$y' x \frac{-x}{\sqrt{1-x^2}} + \sqrt{1-x^2} x y'' = 0$$

$$\frac{-xy' + (1-x^2)y''}{\sqrt{1-x^2}} = 0$$

$$(1-x^2)y'' - xy' = 0$$

IV.

(a) State the product rule and quotient rule of differentiation

Product rule

$$\frac{d}{dx}(u \cdot v) = u \frac{dv}{dx} + v \frac{du}{dx}$$

Quotient rule

$$\frac{d}{dx}\left(\frac{u}{v}\right) = \frac{v \frac{du}{dx} - u \frac{dv}{dx}}{v^2}$$

(b) Find  $\frac{dy}{dx}$ , if

i.  $Y = \frac{1+\cos x}{(x+\sin x)^3}$

$$Y = \frac{1+\cos x}{(x+\sin x)^3}$$

$$\frac{dy}{dx} = \frac{(x+\sin x)^3 x - \sin x - (1+\cos x) \times 3(x+\sin x)^2 (1+\cos x)}{(x+\sin x)^4}$$

ii.  $Y = \log(x + \sqrt{1+x^2})$

$$Y = \log(x + \sqrt{1+x^2})$$

$$\frac{dy}{dx} = \frac{1}{x + \sqrt{1+x^2}} \times \left(1 + \frac{1}{2\sqrt{1+x^2}} \cdot 2x\right)$$

iii.  $Y = \sec^2(e^x)$

$$Y = \sec^2(e^x)$$

$$\frac{dy}{dx} = 2\sec(e^x) \times \sec(e^x)\tan(e^x) \times e^x$$

(c) If  $x = a(\theta + \sin\theta) = a\theta + a\sin\theta$

$$y = a(1 - \cos\theta) = a - a\cos\theta \quad \text{prove that } \frac{dy}{dx} = \tan \frac{\theta}{2}$$

$$\frac{dx}{d\theta} = a + a\cos\theta$$

$$\frac{dy}{d\theta} = a\sin\theta$$

$$\therefore \frac{dx}{dy} = \frac{\frac{dy}{d\theta}}{\frac{dx}{d\theta}} = \frac{a\sin\theta}{a + a\cos\theta}$$

$$= \frac{a\sin\theta}{a(1 + \cos\theta)}$$

$$= \frac{\sin\theta}{2\cos^2\left(\frac{\theta}{2}\right)}$$

$$= \frac{2\sin\left(\frac{\theta}{2}\right)\cos\left(\frac{\theta}{2}\right)}{2\cos^2\left(\frac{\theta}{2}\right)}$$

$$= \frac{\sin\left(\frac{\theta}{2}\right)}{\cos\left(\frac{\theta}{2}\right)}$$

$$= \tan\left(\frac{\theta}{2}\right)$$

$$\frac{dy}{dx} = \tan \frac{\theta}{2}$$

V.

(a) Find the equation of a tangent and normal to the curve  $y = 2\log x$  at the point  $(1, 0)$

$$Y = 2\log x$$

$$\frac{dy}{dx} = \frac{2}{x}$$

$$\therefore \frac{dy}{dx} \text{ at } (1, 0) = \frac{2}{1} = 2$$

$$\therefore m = 2$$

$$X_1 = 1$$

$$Y_1 = 0$$

$$M = 2$$

Equation of tangent is  $y - y_1 = m(x - x_1)$

$$Y - 0 = 2(x - 1)$$

$$Y = 2x - 2$$

Equation of normal is  $y - y_1 = -\frac{1}{m}(x - x_1)$

$$Y - 0 = -\frac{1}{2}(x - 1)$$

$$Y = -\frac{1}{2}(x - 1)$$

$$Y = -\frac{1}{2}x + \frac{1}{2}$$

- (b) A circular path of oil spreads out on water at the rate of  $12 \text{ cm}^2/\text{min}$ . how fast is the radius increasing when the radius is 2cms

Area of the circle =  $\pi r^2$

$$A = \pi r^2$$

$$\therefore \frac{dA}{dt} = \pi \times 2r \times \frac{dr}{dt}$$

But given  $\frac{dA}{dt} = 12 \text{ cm}^2/\text{min}$

$$12 = \pi \times 2r \times \frac{dr}{dt}$$

$$12 = \pi \times 2 \times 2 \times \frac{dr}{dt}$$

$$12 = \pi \times 4 \times \frac{dr}{dt}$$

$$\frac{12}{4\pi} = \frac{dr}{dt}$$

$$\frac{dr}{dt} = \frac{3}{\pi} \text{ cm/min}$$


- (c) An open box is to be made out a square sheet of side 18cms. By cutting off equal shares at each and turning up the sides. What size of the squares should be cut in order that the volume of the box may be maximum?

Length of the box =  $18 - 2x$

Breadth of the box =  $18 - 2x$

Height =  $x$

$$\begin{aligned} \therefore \text{Volume } v &= (18 - 2x)(18 - 2x)x \\ &= (324 - 36x - 36x + 4x^2) x \\ &= 324x - 72x^2 + 4x^3 \end{aligned}$$



$$V = 4x^3 - 72x^2 + 324x$$

$$\frac{dv}{dx} = 0$$

$$12x^2 - 72x + 324 = 0$$

$$12(x^2 - 12x + 27) = 0$$

At a maxima or minima

$$\frac{dv}{dx} = 0 \implies 12(x^2 - 12x + 27) = 0$$

$$x^2 - 12x + 27 = 0$$

$$x(x - 9)(x - 3) = 0$$

$$x = 3, \text{ or } 9$$

But  $x = 9$  is not possible

$$\therefore x = 3$$

VI.

- (a) If  $q = \frac{10p}{p-10}$ , find the rate of change of  $q$  change with respect to 'p'

$$\frac{dq}{dp} = \frac{(p-10) \times 10 - 10p \times 1}{(p-10)^2}$$

$$\frac{dq}{dp} = \frac{10p - 100 - 10p}{(p-10)^2}$$

$$\frac{dq}{dp} = \frac{-100}{(p-10)^2}$$

- (b) Find the equation of the tangent to the curve  $x = \cos 2t$ ,  $y = \sin 2t$  at  $t = \pi/8$

$$\frac{dy}{dx} = \frac{\frac{dy}{dt}}{\frac{dx}{dt}}$$

$$= \frac{2\cos 2t}{-2\sin 2t}$$

$$= -\cot 2t$$

$$\therefore m = -\cot 2\frac{\pi}{8} = -1$$

Equation of tangent is

$$y - y_1 = m(x - x_1)$$

$$(x_1, y_1) = \left(\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}\right)$$

$$y - \frac{1}{\sqrt{2}} = -1\left(x - \frac{1}{\sqrt{2}}\right)$$

$$M = -1$$

$$y - \frac{1}{\sqrt{2}} = -x + \frac{1}{\sqrt{2}}$$

- (c) The displacement of a particle at time 't' is  $s = \frac{1}{2}t^2 + \sqrt{t}$ . Find the velocity at time  $t = 9$  seconds.

$$s = \frac{1}{2}t^2 + \sqrt{t}$$

$$\text{Velocity} = \frac{ds}{dt} = \frac{1}{2} \cdot 2t + \frac{1}{2\sqrt{t}}$$

$$\frac{ds}{dt} = t + \frac{1}{2\sqrt{t}}$$

$$\text{At } t = 9 \text{ secs, } \frac{ds}{dt} = 9 + \frac{1}{2\sqrt{9}}$$

$$= 9 + \frac{1}{2 \times 3}$$

$$= 9 + \frac{1}{6}$$

$$\frac{ds}{dt} = \frac{55}{6}$$

- (d) A hollow cylindrical vessel to hold 100cc of water is to be made. So that the area of the metal

used is minimum. Prove that the radius which will give minimum area is  $\sqrt[3]{\left(\frac{100}{\pi}\right)}$  cms

$$V = \pi r^2 h$$

$$100 = \pi r^2 h$$

$$\therefore h = \frac{100}{\pi r^2} \text{ (1)}$$

$$\text{Surface area } S = \pi r^2 + 2\pi r h$$

From (1),

$$S = \pi r^2 + 2\pi r \left(\frac{100}{\pi r^2}\right)$$

$$S = \pi r^2 + \frac{200}{r} \quad (2)$$

If the surface area y minimum

$$1) \frac{ds}{dr} = 0$$

$$2) \frac{d^2s}{dr^2} > 0$$

$$\frac{ds}{dr} = 0 \implies \frac{d}{dr} \left( \pi r^2 + \frac{200}{r^2} \right) = 0$$

$$\implies 2\pi r - \frac{200}{r^2} = 0$$

$$\implies 2\pi r = \frac{200}{r^2}$$

$$\implies r^3 = \frac{200}{2\pi}$$

$$\implies r^3 = \frac{100}{\pi}$$

$$\therefore r = \sqrt[3]{\left(\frac{100}{\pi}\right)}$$

Check whether the surface area is minimum, at this value

$$\frac{d^2s}{dr^2} = \frac{d}{dr} \left( 2\pi r + \frac{200}{r^2} \right)$$

$$= 2\pi r + \frac{400}{r^3} > 0$$

$\therefore$  The radius which will give minimum surface area is  $\sqrt[3]{\left(\frac{100}{\pi}\right)} \text{cm}$

VII.

$$(a) \int \frac{x^2}{1+x^6} dx$$

$$\text{Put } u = x^3$$

$$\frac{du}{dx} = 3x^2$$

$$\frac{du}{3} = x^2 dx$$

$$\begin{aligned}\int \frac{x^2}{1+x^6} dx &= \int \frac{\frac{du}{3}}{1+u^2} dx \\ &= \frac{1}{3} \int \frac{1}{1+u^2} dx \\ &= \frac{1}{3} \tan^{-1} u \\ &= \frac{1}{3} \tan^{-1} x^3 + c\end{aligned}$$

$$(b) \int \frac{1}{1-7x}$$

$$\int \frac{1}{1-7x} = \frac{\log(1-7x)}{-7} + c$$

$$(c) \int_{\frac{\pi}{4}}^{\frac{\pi}{2}} \frac{\sin 2x}{\sin^2 x} dx$$

$$\begin{aligned}\int_{\frac{\pi}{4}}^{\frac{\pi}{2}} \frac{\sin 2x}{\sin^2 x} dx &= \int_{\frac{\pi}{4}}^{\frac{\pi}{2}} \frac{2 \sin x \cos x}{\sin^2 x} dx \\ &= \int_{\frac{\pi}{4}}^{\frac{\pi}{2}} 2 \frac{\cos x}{\sin x} dx \\ &= 2 \int_{\frac{\pi}{4}}^{\frac{\pi}{2}} \cot x dx \\ &= 2 [\log \sin x]_{\frac{\pi}{4}}^{\frac{\pi}{2}} \\ &= 2 [\log \sin(\frac{\pi}{2}) - \log \sin(\frac{\pi}{4})] \\ &= 2 [\log 1 - \log \frac{1}{\sqrt{2}}] \\ &= 2 [0 - \log \frac{1}{\sqrt{2}}] \\ &= 2 [-[(\log 1) - \log \sqrt{2}]] \\ &= 2 [-(0 - \log \sqrt{2})]\end{aligned}$$

$$= 2[\log\sqrt{2}]$$

$$= \log(\sqrt{2})^2$$

$$= \log 2$$

$$(d) \int x^2 e^{x^3} dx$$

$$U = x^3$$

$$\frac{du}{dx} = 3x^2$$

$$x^2 dx = \frac{du}{3}$$

$$\begin{aligned} \therefore \int x^2 e^{x^3} dx &= \int \frac{du}{3} e^u \\ &= \frac{1}{3} \int e^u du \\ &= \frac{1}{3} e^u \\ &= \frac{1}{3} e^{x^3} + c \end{aligned}$$

$$(e) \int \sin^{-1} x dx$$

$$\int \sin^{-1} x dx = \int \sin^{-1} x \times 1 dx$$

$$= \sin^{-1} x \times x - \int x \times \frac{1}{\sqrt{1-x^2}} dx$$

$$= x \sin^{-1} x - \int \frac{x}{\sqrt{1-x^2}} dx \quad \text{---(1)}$$

$$\text{Put } u = 1 - x^2$$

$$\frac{du}{dx} = -2x$$

$$\frac{du}{-2} = x dx$$

$$\int \sin^{-1} x dx = x \sin^{-1} x - \int \frac{-du}{\sqrt{u}}$$

$$= x \sin^{-1} x + \int \frac{1}{2} \cdot \frac{1}{\sqrt{u}} du$$

$$= x \sin^{-1} x + \sqrt{u}$$

$$= x \sin^{-1} x + \sqrt{1-x^2}$$



VIII.

(a) Find  $\int \frac{4x+2}{x^2+x+3} dx$

$$\begin{aligned}\int \frac{4x+2}{x^2+x+3} dx &= \int \frac{2(2x+1)}{(x^2+x+3)} dx \\ &= \int \frac{2du}{u} dx \\ &= 2 \int \frac{1}{u} du \\ &= 2 \log u \\ &= 2 \log(x^2 + x + 3) \\ &= \log(x^2 + x + 3)^2\end{aligned}$$

$$u = x^2 + x + 3$$

$$\frac{du}{dx} = 2x + 1$$

$$du = (2x + 1) dx$$

(b)  $\int \frac{\tan^{-1} 5x}{1+25x^2} dx$

$$= \int u \frac{du}{5}$$

$$= \frac{1}{5} \frac{u^2}{2}$$

$$= \frac{1}{5} \frac{(\tan^{-1} 5x)^2}{2} + c$$

$$u = \tan^{-1} 5x$$

$$\frac{du}{dx} = \frac{1}{1+25x^2} \times 5$$

$$\frac{du}{5} = \frac{1}{1+25x^2} dx$$

(c)  $\int \sqrt{4x-3} dx$

$$= \int (4x-3)^{\frac{1}{2}} dx$$

$$= \frac{(4x-3)^{\frac{3}{2}}}{\frac{1}{2} \times 4}$$

=

$$= (4x-3)^{\frac{3}{2}} \times \frac{1}{6} + c$$

$$(d) \int_0^1 x^2(x+1)^2 dx$$

$$\begin{aligned}\int_0^1 x^2(x+1)^2 dx &= \int_0^1 x^2(x^2 + 2x + 1) dx \\ &= \int_0^1 (x^4 + 2x^3 + x^2) dx \\ &= \left[ \frac{x^5}{5} + \frac{2x^4}{4} + \frac{x^3}{3} \right]_0^1 \\ &= \frac{1}{5} + \frac{2}{4} + \frac{1}{3} \\ &= \frac{1}{5} + \frac{1}{2} + \frac{1}{3} \\ &= \frac{1}{5} + \frac{5}{6} \\ &= \frac{6+25}{30} = \frac{31}{30}\end{aligned}$$

$$(e) \int x \cos x dx$$

$$\begin{aligned}\int x \cos x dx &= x \sin x - \int \sin x \times 1 dx \\ &= x \sin x + \int \sin x dx \\ &= x \sin x + \cos x + c\end{aligned}$$

IX.

(a) Find the area enclosed between one arch of the curve  $\sin 3x$  and the x axis

$$Y = \sin 3x \quad x = 0, \frac{\pi}{3}$$

$$\therefore \text{Area } A = \int_a^b y dx$$

$$A = \int_0^{\frac{\pi}{3}} \sin 3x dx$$

$$= \left[ -\frac{\cos 3x}{3} \right]_0^{\frac{\pi}{3}}$$

$$= \left( -\frac{\cos \pi}{3} + \frac{\cos 0}{3} \right)$$

$$= \left( -\frac{-1}{3} + \frac{1}{3} \right)$$

$$= \left(\frac{1}{3} + \frac{1}{3}\right)$$

$$= \left(\frac{2}{3}\right) \text{square units}$$

(b) Show by integration, the volume of a right circular cone of height  $h$  & base radius is  $\frac{1}{3}\pi r^2 h$

$$Y = mx$$

$$Y = \tan \theta x$$

$$Y = \frac{r}{h} x$$

$$\therefore V = \pi \int_0^h y^2 dx$$


$$= \pi \int_0^h \left(\frac{r}{h} x\right)^2 dx$$

$$= \pi \int_0^h \frac{r^2}{h^2} x^2 dx$$

$$= \frac{\pi r^2}{h^2} \left[\frac{x^3}{3}\right]_0^h$$

$$= \frac{\pi r^2}{h^2} \frac{h^3}{3}$$

$$= \frac{1}{3} \pi r^2 h$$



(c) Solve  $x \frac{dy}{dx} + y = x^2$

Dividing both sides by  $x$

$$\frac{dy}{dx} + \frac{y}{x} = \frac{x^2}{x}$$

$$\frac{dy}{dx} + \frac{1}{x}y = x$$

This is the form  $\frac{dy}{dx} + Py = Q$

$$P = \frac{1}{x} \quad Q = x$$

$$\text{I.F} = e^{\int P dx}$$

$$= e^{\int \frac{1}{x} dx} = e^{\log x} = x$$

∴ Solution is

$$Y \times \text{IF} = \int (Q \times \text{IF}) dx$$

$$Y \times x = \int (x \times x) dx$$

$$Yx = \int x^2 dx$$

$$xy = \frac{x^3}{3} + c$$

X.

(a) Find the area bounded by the curve  $y = x^2$  and  $y = 3x$

$$f(x) = x^2$$

$$g(x) = 3x$$

$$x^2 = 3x$$

$$x^2 - 3x = 0$$

$$x(x - 3) = 0$$

$$x = 0 \text{ or } x = 3$$

$$\therefore \text{Area} = \int_a^b f(x) - g(x) dx$$

$$= \int_0^3 (x^2 - 3x) dx$$

$$= \left[ \frac{x^3}{3} - \frac{3x^2}{2} \right]_0^3$$

$$\text{Area} = \frac{3^3}{3} - \frac{3 \times 3^2}{2} - \left( \frac{0^3}{3} - \frac{3 \times 0^2}{2} \right)$$

$$= \frac{27}{3} - \frac{27}{2} - 0$$

$$= \frac{54 - 81}{6}$$

$$= -\frac{27}{6}$$

$$= -\frac{9}{2}$$

$$\text{Area} = \frac{9}{2} \text{square units (area cannot -ve)}$$

(b) Find the volume of a sphere of radius r

$$\text{We have } V = \pi \int_a^b y^2 dx$$

$\therefore$  Volume of the solid generated by rotation of Quadrant OAB

$$\begin{aligned} &= \pi \int_a^b y^2 dx \\ &= \pi \int_a^b r^2 - x^2 dx \\ &= \pi \int_0^r r^2 - x^2 dx \\ &= \pi \left[ r^2 x - \frac{x^3}{3} \right]_0^r \\ &= \pi \left[ r^3 - \frac{r^3}{3} \right] \\ &= \pi \left[ \frac{2r^3}{3} \right] \\ &= \left[ \frac{2\pi r^3}{3} \right] \\ &= \frac{2}{3} \pi r^3 \end{aligned}$$


$\therefore$  Volume of sphere = 2 x volume of OAB

$$\begin{aligned} &= 2 \times \frac{2}{3} \pi r^3 \\ &= \frac{4}{3} \pi r^3 \end{aligned}$$

(c) solve  $\frac{dy}{dx} + \sqrt{\frac{1-y^2}{1-x^2}} = 0$

$$\frac{dy}{dx} + \sqrt{\frac{1-y^2}{1-x^2}} = 0$$

$$\frac{dy}{dx} = -\sqrt{\frac{1-y^2}{1-x^2}}$$



$$\frac{dy}{\sqrt{1-y^2}} = -\frac{dx}{\sqrt{1-x^2}}$$

Integrating on both sides we get

$$\int \frac{dy}{\sqrt{1-y^2}} = - \int \frac{dx}{\sqrt{1-x^2}}$$

$$\sin^{-1} y = \sin^{-1} x$$

$$\sin^{-1} y + \sin^{-1} x = c$$

www.madimpoly.com