

SECOND SEMESTER DIPLOMA EXAMINATION IN ENGINEERING/  
TECHNOLIGY- OCTOBER, 2013

TECHNICAL MATHEMATICS- II  
(Common – Except DCP and CABM)

[Time: 3 hours

(Maximum marks: 100)

Marks

PART –A  
(Maximum marks: 10)

(Answer all questions. Each question carries 2 marks)

I.

(a) Find the derivatives of  $y = 3\cos x - 4\tan x$

$$y = 3\cos x - 4\tan x$$

$$\begin{aligned}\frac{dy}{dx} &= 3 \times -\sin x - 4\sec^2 x \\ &= -3\sin x - 4\sec^2 x\end{aligned}$$

(b) Evaluate  $\lim_{x \rightarrow 0} \frac{2x-3}{3x+4}$

$$\lim_{x \rightarrow 0} \frac{2x-3}{3x+4} = \frac{2 \times 0 - 3}{3 \times 0 + 4} = \frac{-3}{4}$$

(c) Check whether the function  $x^2 - 3x + 2$  is decreasing at  $x = 1$

$$y = x^2 - 3x + 2$$

$$\frac{dy}{dx} = 2x - 3$$

At  $x = 1$

$$\frac{dy}{dx} = 2 \times 1 - 3 = 2 - 3 = -1 < 0$$

$$\therefore \frac{dy}{dx} < 0 \text{ at } x = 1$$

$\therefore$  Function is decreasing at  $x = 1$

(d) Find  $\int (2x + 1)^2 dx$

$$\begin{aligned}\int (2x + 1)^2 dx &= \int 4x^2 + 1 + 4x dx \\ &= 4 \int x^2 dx + \int 1 dx + \int 4x dx \\ &= 4 \frac{x^3}{3} + x + 4 \frac{x^2}{2} + C\end{aligned}$$

(e) Solve  $\frac{d^2y}{dx^2} = \sec^2 x$

$$\frac{d}{dx} \left( \frac{dy}{dx} \right) = \sec^2 x$$

Integrating on both sides we get,

$$\frac{dy}{dx} = \tan x + C_1$$

Again integrating on both sides we get,

$$y = \log \sec x + C_1 x + C_2$$

## PART -B

Answer any five questions. Each question carries 6 marks

II.

(a)

i. Find the differential coefficient of 'tanx' using quotient rule.

$$Y = \tan x = \frac{\sin x}{\cos x}$$

We know by quotient rule,

$$\frac{dy}{dx} = \frac{d}{dx} \left( \frac{u}{v} \right) = \frac{v \times \frac{du}{dx} - u \times \frac{dv}{dx}}{v^2}$$

$$\frac{d}{dx} (\tan x) = \frac{\cos x \times \frac{d(\sin x)}{dx} - \sin x \times \frac{d(\cos x)}{dx}}{\cos^2 x}$$

$$= \frac{\cos x \times \cos x - \sin x \times -\sin x}{\cos^2 x}$$

$$= \frac{\cos^2 x + \sin^2 x}{\cos^2 x}$$

$$= \frac{1}{\cos^2 x} = \sec^2 x$$

ii. Evaluate  $\lim_{x \rightarrow \infty} \frac{x^2 - 2x + 8}{4x^3 - 3}$

$$\lim_{x \rightarrow \infty} \frac{x^2 - 2x + 8}{4x^3 - 3}$$

$$= \lim_{x \rightarrow \infty} \frac{x^2 \left[ 1 - \frac{2x}{x^2} + \frac{8}{x^3} \right]}{x^3 \left[ 4 - \frac{3}{x^3} \right]}$$

$$= \lim_{x \rightarrow \infty} \frac{1 \left[ 1 - \frac{2x}{x^2} + \frac{8}{x^3} \right]}{x \left[ 4 - \frac{3}{x^3} \right]}$$

$$\left[ \lim_{x \rightarrow \infty} \frac{1}{x} = 0 \right]$$

$$= 0 \frac{\left( 1 - \frac{2}{\infty} + \frac{8}{\infty} \right)}{\left( 4 - \frac{3}{\infty} \right)}$$

$$= 0$$

(b) If  $y = \sin^{-1} x$ , show that  $(1 - x^2) \frac{d^2 y}{dx^2} - x \frac{dy}{dx} = 0$

$$y = \sin^{-1} x$$

$$\frac{dy}{dx} = \frac{1}{\sqrt{1-x^2}}$$

$$\frac{dy}{dx} \sqrt{1-x^2} = 1$$

Differentiating on both sides,

$$\frac{dy}{dx} \times \frac{1}{2\sqrt{1-x^2}} \times -2x + \sqrt{1-x^2} \frac{d^2y}{dx^2} = 0$$

$$\frac{-x dy}{dx} \times \frac{1}{\sqrt{1-x^2}} + \sqrt{1-x^2} \frac{d^2y}{dx^2} = 0$$

Multiplication on both sides by  $\sqrt{1-x^2}$  we get,

$$\frac{-x dy}{dx} + (1-x^2) \frac{d^2y}{dx^2} = 0$$

$$\text{Or } (1-x^2) \frac{d^2y}{dx^2} - \frac{x dy}{dx} = 0$$

(c) For what value of  $x$  tangent to the curve  $y = \frac{x}{(1-x)^2}$  will be parallel to:

- i. X axis
- ii. Y axis
- i. Parallel to x axis

$$Y = \frac{x}{(1-x)^2}$$

$$\frac{dy}{dx} = \frac{(1-x)^2 \times 1 - x \times 2(1-x) \times -1}{((1-x)^2)^2}$$

If  $y = \frac{x}{(1-x)^2}$  is parallel to x axis then,

$$\frac{dy}{dx} = 0 \implies \frac{(1-x)^2 + 2x(1-x)}{(1-x)^4}$$

$$\frac{1-x^2}{(1-x)^4} = 0$$

$$\frac{(1+x)(1-x)}{(1-x)^4} = 0$$

$$\frac{(1-x)}{(1-x)^3} = 0$$

$$x = -1$$

ii. Parallel to y axis

If  $y = \frac{x}{(1-x)^2}$  is parallel to y axis then

$$\text{Slop} = \infty$$

$$\text{Ie, } \frac{dy}{dx} = \infty \implies \frac{(1-x)^2 + 2x(1-x)}{(1-x)^4} = 0$$

$$\implies (1-x)^4 = 0$$

$$\implies x = 1$$

(d) Find the minimum value of  $y = 4x^3 + 9x^2 - 12x + 2$

$$y = 4x^3 + 9x^2 - 12x + 2$$

$$\frac{dy}{dx} = 12x^2 + 18x - 12 = 0$$

$$\frac{dy}{dx} = 0 \implies 12x^2 + 18x - 12 = 0$$

$$6(2x^2 + 3x - 2) = 0$$

$$x = \frac{-3 \pm \sqrt{9+16}}{4} = \frac{-3 \pm 5}{4}$$

$$= -2, \frac{1}{2}$$

$$\frac{d^2y}{dx^2} = 24x + 18$$

At minimum

$$\frac{dy}{dx} = 0 \& \frac{d^2y}{dx^2} > 0$$

At  $x = -2$

$$\frac{d^2y}{dx^2} = 24x - 2 + 18 = -48 + 18 = -3 < 0$$

At  $x = \frac{1}{2}$

$$\frac{d^2y}{dx^2} = 24x^{1/2} + 18 = 12 + 18 = 30 > 0$$

At  $x = 1/2$   $y$  is minimum

$$\begin{aligned} \therefore \text{Minimum value } y &= 4x \left(\frac{1}{2}\right)^3 + 9\left(\frac{1}{2}\right)^2 - 12x \frac{1}{2} + 2 \\ &= -\frac{5}{4} \end{aligned}$$

(e) Find

i.  $\int \frac{x^2+3x-2}{x} dx$

ii.  $\int \sin 3x \cos x dx$

$$\begin{aligned} \text{i. } \int \frac{x^2+3x-2}{x} dx &= \int \frac{x^2}{x} dx + \int \frac{3x}{x} dx - \int \frac{2}{x} dx \\ &= \int x dx + \int 3 dx - 2 \int \frac{1}{x} dx \\ &= \frac{x^2}{2} + 3x - 2 \log x + C \end{aligned}$$

ii.  $\sin A \cos B = \frac{1}{2} [\sin(A+B) + \sin(A-B)]$

$$\sin 3x \cos x = \frac{1}{2} [\sin 4x + \sin 2x]$$

$$\int \sin 3x \cos x dx = \frac{1}{2} \int (\sin 4x + \sin 2x) dx$$

$$= \frac{1}{2} \int \sin 4x dx + \frac{1}{2} \int \sin 2x dx$$

$$= \frac{1}{2} \left( \frac{-\cos 4x}{4} \right) + \frac{1}{2} \left( \frac{-\cos 2x}{2} \right)$$

$$= \frac{-\cos 4x}{8} + \frac{-\cos 2x}{4} + C$$

(f) Find  $\int \tan^{-1} x \, dx$

$$\begin{aligned}\text{We have } \int \tan^{-1} x \, dx &= \int \tan^{-1} x \times 1 \, dx \\ &= \tan^{-1} x \int 1 \, dx - \int \frac{d}{dx} (\tan^{-1} x) \int 1 \, dx \, dx \\ &= \tan^{-1} x \times x - \int \frac{1}{1+x^2} x \, dx \\ &= x \tan^{-1} x - \frac{1}{2} \log(1+x^2) + C\end{aligned}$$

(g) Solve  $\frac{dy}{dx} = \frac{xy^2+x}{yx^2+x}$

$$\frac{dy}{dx} = \frac{x(y^2+1)}{y(x^2+1)}$$

$$dy \cdot y(x^2+1) = dx \cdot x(y^2+1)$$

$$\frac{ydy}{y^2+1} = \frac{xdx}{x^2+1} \quad \text{Integrating both sides}$$

$$\int \frac{ydy}{y^2+1} = \int \frac{xdx}{x^2+1} \quad (1)$$

$$\text{Consider } \int \frac{x}{x^2+1} dx \quad \text{put } u = x^2 + 1$$

$$\frac{du}{dx} = 2x \quad du = 2x \, dx \quad \frac{du}{2} = x \, dx$$

$$\int \frac{x}{x^2+1} dx = \int \frac{du}{2u} = \frac{1}{2} \int \frac{du}{u} = \frac{1}{2} \log u$$

$$= \frac{1}{2} \log(x^2 + 1) + C_1$$

Similarly  $\int \frac{y dy}{y^2+1} = \log \frac{(y^2+1)}{2} + C_2$

(1) Becomes  $\frac{1}{2} \log(y^2 + 1) + C_2 = \frac{1}{2} \log(x^2 + 1) + C_1$

$$\frac{1}{2} \log(y^2 + 1) - \frac{1}{2} \log(x^2 + 1) + C_1 = C \quad [\text{take } C_1 - C_2 = C]$$

$$\implies \frac{1}{2} \log \frac{1+y^2}{1+x^2} = C$$

$$\implies \log \frac{1+y^2}{1+x^2} = 2C$$

$$\implies 1 + y^2 = e^{2c}(1 + x^2)$$

$$\implies 1 + y^2 = k(1 + x^2)$$

PART -C

(Maximum mark: 60)

Answer four full questions. Each question carries 15 marks.

III.

(a) Using 1<sup>st</sup> principles, find the derivative of sinx

By 1<sup>st</sup> principle we have

$$\frac{dy}{dx} = \lim_{\Delta x \rightarrow 0} \frac{f(x+\Delta x) - f(x)}{\Delta x} \quad \text{_____ (1)}$$

$$f(x) = \sin x$$

$$f(x + \Delta x) = \sin(x + \Delta x)$$

Then (1) becomes

$$\begin{aligned} \frac{dy}{dx} &= \lim_{\Delta x \rightarrow 0} \frac{\sin(x+\Delta x) - \sin x}{\Delta x} \\ &= \lim_{\Delta x \rightarrow 0} \frac{2 \left[ \frac{\cos(2x+\Delta x)}{2} \sin\left(\frac{\Delta x}{2}\right) \right]}{\Delta x} \end{aligned}$$



$$= \lim_{\Delta x \rightarrow 0} \frac{2 \left[ \frac{\cos(2x+\Delta x)}{2} \sin\left(\frac{\Delta x}{2}\right) \right]}{\frac{\Delta x}{2} \cdot 2}$$

$$= \cos\left(\frac{2x+0}{2}\right) = \cos x$$

(b) Find  $\frac{dy}{dx}$  if:

i.  $y = e^x \tan x$

$$y = e^x \tan x \qquad \left[ \frac{d}{dx}(u \cdot v) = u \frac{dv}{dx} + v \frac{du}{dx} \right]$$

$$\frac{dy}{dx} = e^x \sec^2 x + \tan x e^x$$

ii.  $y = \log(\sec x + \tan x)$

$$\begin{aligned} \frac{dy}{dx} &= \frac{dy}{dx} \log(\sec x + \tan x) \\ &= \frac{1}{\sec x + \tan x} (\sec x \tan x + \sec^2 x) \\ &= \frac{\sec x (\tan x + \sec x)}{(\sec x + \tan x)} = \sec x \end{aligned}$$

(c) If  $ax^2 + 2hxy + by^2 = 0$  find  $\frac{dy}{dx}$

$$ax^2 + 2hxy + by^2 = 0$$

Differentiating on both sides

$$2ax + 2h \left( x \frac{dy}{dx} + y \times 1 \right) + 2by \frac{dy}{dx} = 0$$

$$\frac{dy}{dx} (2hx + 2by) + 2ax + 2hy = 0$$

$$\therefore \frac{dy}{dx} = \frac{-(ax+hy)}{(hx+by)}$$

IV.

(a) If  $x = a(\theta + \sin\theta)$ ,  $y = a(1 + \cos\theta)$ , find  $\frac{dy}{dx}$

$$\frac{dy}{dx} = \frac{\frac{dy}{d\theta}}{\frac{dx}{d\theta}}$$

$$\frac{dy}{d\theta} = a(-\sin\theta) = -a \sin\theta = -2a \sin\left(\frac{\theta}{2}\right) \cos\left(\frac{\theta}{2}\right)$$

$$\frac{dx}{d\theta} = a(1 + \cos\theta) = 2a \cos^2\left(\frac{\theta}{2}\right)$$

(1) Becomes

$$\begin{aligned} \frac{dy}{dx} &= \frac{-2a \sin\left(\frac{\theta}{2}\right) \cos\left(\frac{\theta}{2}\right)}{2a \cos^2\left(\frac{\theta}{2}\right)} \\ &= -\tan\left(\frac{\theta}{2}\right) \end{aligned}$$

(b) If  $y = ae^x + be^{2x}$  show that  $y'' - 3y' + 2y = 0$

$$y = ae^x + be^{2x}$$

$$y' = ae^x + 2be^{2x}$$

$$y'' = ae^x + 4be^{2x}$$

$$\begin{aligned} y'' - 3y' + 2y &= (ae^x + 4be^{2x}) - 3(ae^x + 2be^{2x}) + 2(ae^x + be^{2x}) \\ &= ae^x + 4be^{2x} - 3ae^x - 6be^{2x} + 2ae^x + 2be^{2x} \\ &= 0 \end{aligned}$$

(c) Find  $\frac{d^2y}{dx^2}$ , if  $y = \sin x \cos x$

$$y = \sin x \cos x$$

$$\frac{dy}{dx} = \sin x \times \frac{d}{dx} \cos x + \cos x \times \frac{d}{dx} \sin x$$

$$= \sin x \times (-\sin x) + \cos x \times \cos x$$

$$= -\sin^2 x + \cos^2 x$$

$$= \cos^2 x - \sin^2 x$$

$$= \cos 2x$$

V.

- (a) If the displacement of a body is given by  $s = 2t^3 - 3t^2 - 12t + 6$ , find when the body attains the greatest height and also find the acceleration then.

$$s = 2t^3 - 3t^2 - 12t + 6$$

At maximum

$$\frac{ds}{dt} = 0, \frac{d^2s}{dt^2} < 0$$

$$\frac{ds}{dt} = 0 \implies 6t^2 - 6t - 12 = 0$$

$$6(t^2 - t - 2) = 0$$

$$t = \frac{1 \pm \sqrt{1+8}}{2} = \frac{1 \pm 3}{2} = 2, -1$$

$$\frac{d^2s}{dt^2} = 12t - 6$$

At  $t = 2$

$$\begin{aligned} \frac{d^2s}{dt^2} &= 12 \times 2 - 6 \\ &= 24 - 6 = 18 > 0 \end{aligned}$$

At  $t = -1$

$$\begin{aligned} \frac{d^2s}{dt^2} &= 12 \times -1 - 6 \\ &= -12 - 6 = -18 < 0 \end{aligned}$$

At  $t = -1$

$$\begin{aligned} \text{Height } s &= 2(-1)^3 - 3(-1)^2 - 12 \times -1 + 6 \\ &= -2 - 3 + 12 + 6 = 13 \end{aligned}$$

$$\text{Acceleration is } \frac{d^2s}{dt^2} = -18$$

- (b) A stone is dropped into still water. The radius of the outermost ripple then formed increases at the rate of 6cm/sec. how fast is the area increasing when the radius is 16cms.

$$\text{Area of circle } A = \pi r^2$$

$$\frac{dA}{dt} = 2\pi r \frac{dr}{dt} \quad \text{_____ (1)}$$

$$\frac{dr}{dt} = 6 \text{ cm/sec}$$

Then (1) becomes

$$\frac{dA}{dt} = 2\pi \times 16 \times 6$$

$$= 2\pi \times 96$$

$$= 192\pi \text{ cm}^2/\text{sec}$$

$\therefore$  Area is increasing at the rate of  $192\pi \text{ cm}^2/\text{sec}$  when radius is 16cm.

- (c) Show that a rectangle of fixed perimeter has its maximum area when it becomes a square

Let  $x$  and  $y$  be the length and breadth of the rectangle of fixed perimeter 'p'

$$\text{Ie, } 2x + 2y = p$$

$$2y = p - 2x$$

$$y = \frac{p-2x}{2} \quad \text{_____ (1)}$$

Area of rectangle =  $x \times y$

$$A = x \times y$$

$$= \frac{p-2x}{2} \times x$$

$$= \frac{1}{2}(px - 2x^2)$$

At maximum or minimum  $\frac{dA}{dx} = 0$

$$\frac{dA}{dx} = \frac{d}{dx} \frac{1}{2}(px - 2x^2) = \frac{1}{2}(p - 4x) = 0$$

$$\implies x = \frac{p}{4}$$

$$\therefore y = \frac{p-2x}{2} = \frac{p-2\frac{p}{4}}{2} = \frac{p-\frac{p}{2}}{2} = \frac{p}{4}$$

$$x = \frac{p}{4}$$

$$y = \frac{p}{4}$$

$$\frac{d^2A}{dx^2} = \frac{d}{dx} \frac{1}{2} (p - 4x) = \frac{1}{2} x - 4 < 0$$

Since  $\frac{d^2A}{dx^2} < 0$ , A is maximum at  $x = \frac{p}{4}$ ,  $y = \frac{p}{4}$

VI.

(a) Find the range of values of x for which the function  $x^2 - 3x + 4$  is:

i. Increasing    ii. Decreasing

i.  $y = x^2 - 3x + 4$

$$\frac{dy}{dx} = 2x - 3$$

If the function is increasing  $\frac{dy}{dx} > 0$

I.e.,  $2x - 3 > 0$

$$2x > 3 \implies x > \frac{3}{2}$$

ii. If the function is decreasing  $\frac{dy}{dx} < 0$

$$2x - 3 < 0 \implies 2x < 3$$

$$x < \frac{3}{2}$$

(b) Show that the maximum value of function  $M = 2x^3 - 9x^2 + 12x$  is 5

$$M = 2x^3 - 9x^2 + 12x$$

At maximum  $\frac{dM}{dx} = 0$  &  $\frac{d^2M}{dx^2} < 0$

$$\frac{dM}{dx} = 0 \implies 6x^2 - 18x + 12 = 0$$

$$\implies 6(x^2 - 3x + 2)$$

$$X = \frac{3 \pm \sqrt{9-8}}{2}$$

$$= \frac{3 \pm 1}{2} = 2, 1$$

At  $x = 1$

$$\frac{d^2M}{dx^2} = 12x - 18 = 12 \times 1 - 18 = -6 < 0$$

At  $x = 2$

$$\frac{d^2M}{dx^2} = 12x - 18 = 12 \times 2 - 18 = 24 - 18 = 6 > 0$$

$\therefore$  M is maximum at  $x = 1$

$$\text{And maximum value } M = 2(1)^3 - 9(1)^2 + 12 \times 1$$

$$= 2 - 9 + 12 = 5$$

- (c) Water is running out of a conical funnel at the rate of 1 cubic inch per second. If the radius of the funnel is 4 inches and altitude is 8 inches. Find the rate at which the water level is dropping when its depth is 6 inches.

Let  $v$  be the volume of the water in the cone.

Let  $\alpha$  be the semi vertical angle of the cone.

Then


$$\tan \alpha = \frac{OA}{OV} = \frac{4}{8} = \frac{1}{2}$$

$$\text{Also } \tan \alpha = \frac{O'A'}{O'V'} = \frac{r}{h}$$

$$\therefore \frac{r}{h} = \frac{1}{2} \implies r = \frac{h}{2}$$

$$\text{Then } V = \frac{1}{3} \pi r^2 h = V = \frac{1}{3} \pi \left(\frac{h}{2}\right)^2 h$$

$$= V = \frac{1}{3} \pi \frac{h^3}{4}$$



$$= \frac{\pi h^3}{12} \text{m}$$

$$\frac{dv}{dt} = -\frac{1}{12} \times \pi \times 3h^2 \frac{dh}{dt}$$

[-ve sign is due to decreasing v]

$$\therefore 1 = -\frac{1}{12} \times \pi \times 3h^2 \frac{dh}{dt} \quad \left[ \text{Given } \frac{dv}{dt} = 1 \right]$$

$$\begin{aligned} \therefore \frac{dh}{dt} &= \frac{-12}{\pi \times 3h^2} = \frac{-4}{\pi h^2} \\ &= \frac{-4}{\pi 6^2} = \frac{-4}{36\pi} \text{inch/sec} \end{aligned}$$

$\therefore$  The water level is dropped at the rate of  $\frac{-4}{36\pi}$  inch/sec

VII. Evaluate

(a)  $\int (2x^3 - 3\sin x + 5x) dx$

$$\begin{aligned} &\int (2x^3 - 3\sin x + 5x) dx \\ &= \int 2x^3 dx - \int 3\sin x dx + \int 5x dx \\ &= \frac{2x^4}{4} - 3x - \cos x + \frac{5x^2}{2} + C \\ &= \frac{x^4}{2} + 3\cos x + \frac{5x^2}{2} + C \end{aligned}$$

(b)  $\int \frac{3x^2}{\sqrt{1-x^6}} dx$

Put  $u = x^3$

$Du = 3x^2 dx$

$$\begin{aligned} \int \frac{3x^2}{\sqrt{1-x^6}} dx &= \int \frac{3x^2}{\sqrt{1-(x^3)^2}} dx \\ &= \int \frac{du}{\sqrt{1-u^2}} = \sin^{-1} u + C \\ &= \sin^{-1} x^3 + C \end{aligned}$$

$$(c) \int x^2 e^x dx$$

[Use ILATE Rule]

Taking  $x^2$  as 1<sup>st</sup> function, we have

$$\begin{aligned} \int x^2 e^x dx &= x^2 \int e^x dx - \int \left( \frac{d}{dx} x^2 \int e^x dx \right) dx \\ &= x^2 e^x - \int 2x \times e^x dx \quad \text{_____ (1)} \end{aligned}$$

$$\begin{aligned} \int x e^x dx &= x \int e^x dx - \int \frac{d}{dx} x \int e^x dx dx \\ &= x e^x - \int e^x dx \\ &= x e^x - e^x + C \quad \text{_____ (2)} \end{aligned}$$

Putting (2) in (1) we have

$$\begin{aligned} \int x^2 e^x dx &= x^2 e^x - 2[x e^x - e^x + C] \\ &= x^2 e^x - 2x e^x + 2e^x + 2C \\ &= x^2 e^x - 2x e^x + 2e^x + C_1 \end{aligned}$$

$$(d) \int_0^{\frac{\pi}{2}} \frac{\cos x}{1 + \sin x} dx \quad \begin{array}{l} \text{Put } u = 1 + \sin x \\ du = \cos x dx \end{array}$$

$$\begin{aligned} \int_0^{\frac{\pi}{2}} \frac{\cos x}{1 + \sin x} dx &= \int_0^{\frac{\pi}{2}} \frac{du}{u} = \log u = \log(1 + \sin x) = f(x) \end{aligned}$$

$$F\left(\frac{\pi}{2}\right) = \log\left(1 + \sin\frac{\pi}{2}\right) = \log 2$$

$$F(0) = \log(1 + \sin 0) = \log 1 = 0$$

$$\int_0^{\frac{\pi}{2}} \frac{\cos x}{1 + \sin x} dx = f\left(\frac{\pi}{2}\right) - f(0) = \log 2$$



VIII. Find

(a)  $\int (4 \sec^2 x + 3 \sin x + e^x) dx$

$$\begin{aligned} & \int (4 \sec^2 x + 3 \sin x + e^x) dx \\ &= 4 \tan x + 3 x - \cos x + e^x + C \\ &= 4 \tan x - 3 \cos x + e^x + C \end{aligned}$$

(b)  $\int_1^e \log x dx$

$$\begin{aligned} \int \log x dx &= \int \log x \times 1 dx \\ &= \log x \times \int 1 dx - \int \frac{d}{dx} \log x \times \int 1 dx dx \\ &= \log x \times x dx - \int \frac{1}{x} x dx \\ &= x \log x - \int 1 dx = x \log x - x \end{aligned}$$

$$\int \log x dx = x \log x - x = f(x)$$

$$f(e) = e \log e - e = 0$$

$$f(1) = 1 \log 1 - 1 = -1$$

$$\therefore \int_1^e \log x dx = f(e) - f(1) = 1$$

(c)  $\int_0^{\frac{\pi}{2}} \cos^2 x dx$

$$\begin{aligned} \int \cos^2 x dx &= \int \frac{1 + \cos 2x}{2} dx \\ &= \int \frac{1}{2} dx + \int \frac{\cos 2x}{2} dx \\ &= \frac{x}{2} + \frac{\sin 2x}{4} = f(x) \end{aligned}$$

$$F\left(\frac{\pi}{2}\right) = \frac{\pi}{2} + \frac{\sin 2\frac{\pi}{2}}{4} = \frac{\pi}{4}$$

$$F(0) = 0$$

$$\therefore \int_0^{\frac{\pi}{2}} \cos^2 x \, dx = F\left(\frac{\pi}{2}\right) - f(0) = \frac{\pi}{4}$$

$$(d) \int (1 + e^{\tan x}) \sec^2 x \, dx$$

Put  $u = \tan x$

$$\int (1 + e^u) \, du = \sec^2 x \, dx$$

$$= \int 1 \, du + \int e^u \, du$$

$$= u + e^u + C$$

$$= \tan x + e^{\tan x} + C$$

IX.

- (a) Find the area enclosed between the curve  $y = x^2 - x + 1$ . The x axis and the ordinate  $x = 1$  and  $x = 3$

$$\text{Area} = \left| \int_a^b f(x) - g(x) \, dx \right|$$

$$f(x) = x^2 - x + 1 \quad g(x) = 0$$

$$a = 1 \quad b = 3$$

$$\text{Required area} = \left| \int_1^3 x^2 - x + 1 \, dx \right|$$

$$\int x^2 - x + 1 \, dx = \int x^2 \, dx - \int x \, dx + \int 1 \, dx$$

$$= \frac{x^3}{3} - \frac{x^2}{2} + x = f(x)$$

$$F(3) = \frac{3^3}{3} - \frac{3^2}{2} + 3 = \frac{27}{3} - \frac{9}{2} + 3$$

$$= \frac{54 - 27}{6} + 34$$

$$= \frac{27}{6} + 3 = \frac{27 + 18}{6}$$

$$= \frac{45}{6}$$

$$F(1) = \frac{1^3}{3} - \frac{1^2}{2} + 1$$

$$= \frac{x^3}{3} - \frac{1}{2} + 1 = \frac{2-3}{6} + 1 = \frac{-1}{6} + 1$$

$$= \frac{-1+6}{6} = \frac{5}{6}$$

Required Area =  $f(3) - f(1)$

$$= \frac{45}{6} - \frac{5}{6} = \frac{20}{3} \text{unit}^2$$

(b) Find the volume of sphere obtained by rotating the circle  $x^2 + y^2 = a^2$  about the x axis

$$\text{is } \frac{4}{3} \pi a^3$$

Equation of circle:  $x^2 + y^2 = a^2$

$$\text{Volume} = \pi \int_{-a}^a y^2 dx$$

$$= \pi \left[ \int_{-a}^a (a^2 - x^2) dx \right]$$

$$= \int (a^2 - x^2) dx = a^2 x - \frac{x^3}{3} = f(x)$$


$$F(a) = a^2 \times a - \frac{a^3}{3} = a^3 - \frac{a^3}{3} = \frac{2a^3}{3}$$

$$F(-a) = a^2 \times -a - \frac{(-a^3)}{3} = -a^3 + \frac{a^3}{3} = -\frac{2a^3}{3}$$

$$F(a) - f(-a) = \frac{4}{3} a^3$$

$$\text{Volume} = \pi \int_{-a}^a y^2 dx = \pi [F(a) - f(-a)]$$

$$= \pi \times \frac{4}{3} a^3$$



(c) Solve  $\frac{dy}{dx} + \sqrt{\left(\frac{1-y^2}{1-x^2}\right)} = 0$

$$\frac{dy}{dx} + \sqrt{\left(\frac{1-y^2}{1-x^2}\right)} = 0$$

$$\frac{dy}{\sqrt{1-y^2}} = \frac{-dx}{\sqrt{1-x^2}}$$

Integrating on both sides,

$$\int \frac{dy}{\sqrt{1-y^2}} = - \int \frac{dx}{\sqrt{1-x^2}}$$

$$\sin^{-1} y = - \sin^{-1} x + C$$


$$\implies \sin^{-1} y + \sin^{-1} x + C$$

X.

(a) Find the area enclosed between one arch of the curve  $y = \sin x$  and the x axis

Required area

$$\begin{aligned} &= \int_a^b f(x) dx \\ &= \int_0^\pi \sin x dx \\ &= [-\cos x]_0^\pi \\ &= [-\cos \pi + \cos 0] \\ &= 2 \text{ unit}^2 \end{aligned}$$



(b) Find the volume of the solid obtained by rotating the area consider the parabola  $y^2 = 4x$  between the ordinates at  $x = 0$ ,  $x = 2$  and the x axis

$$V = \pi \int_a^b y^2 dx$$

$$\begin{aligned} &= \pi \int_0^2 4x \, dx \\ &= \pi \left[ \frac{4x^2}{2} \right]_0^2 \\ &= \pi [2x^2]_0^2 \\ &= \pi [2 \times 2^2] \\ &= 8\pi \text{ cubic unit.} \end{aligned}$$

(c) Solve  $\frac{dy}{dx} + y \cot x = \operatorname{cosec} x$

$$\begin{aligned} \text{Integrating factor} = \text{I.F} &= e^{\int \cot x \, dx} \\ &= e^{\log \sin x} \\ &= \sin x \end{aligned}$$

$$\text{Solution: } y \times \text{IF} = \int \operatorname{cosec} x \times \text{IF} \, dx$$

$$y \sin x = \int dx$$

$$y \sin x = x + C$$